

Bleeding risk scores in atrial fibrillation and venous thromboembolism

Zulkifly, Hanis; Lip, Gregory; Lane, Deirdre

DOI:

[10.1016/j.amjcard.2017.06.058](https://doi.org/10.1016/j.amjcard.2017.06.058)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Zulkifly, H, Lip, G & Lane, D 2017, 'Bleeding risk scores in atrial fibrillation and venous thromboembolism', *The American Journal of Cardiology*. <https://doi.org/10.1016/j.amjcard.2017.06.058>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Eligibility for repository: Checked on 10/7/2017

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Bleeding Risk Scores in Atrial Fibrillation and Venous Thromboembolism

Hanis Zulkifly, MPharm^a, Gregory Yoke Hong Lip, MD^{a, b}, Deirdre Anne Lane, PhD^{a, b}

^aUniversity of Birmingham Institute of Cardiovascular Sciences, City Hospital, Dudley Road, Birmingham B18 7QH, United Kingdom

^bAalborg Thrombosis Research Unit, Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

Corresponding author: Dr Deirdre A. Lane

Telephone number: +44 121-507-5080

Email address: deirdrelane@nhs.net

Address for correspondence: University of Birmingham Institute of Cardiovascular Sciences, City Hospital, Sandwell and West Birmingham Hospitals NHS Trust, Dudley Road, Birmingham B18 7QH, United Kingdom.

Article type: Review article

Running head: Bleeding risk scores in AF and VTE

Funding: None

Abstract

Patients receiving oral anticoagulant (OAC) therapy for stroke prevention in atrial fibrillation (AF) and prevention of venous thromboembolism (VTE) face an increased risk of bleeding with OAC treatment. Clinicians need to weigh-up the benefits of OAC treatment against the risk of bleeding. To help formalise bleeding risk assessment, various bleeding risk scores have been developed to help predict the risk of bleeding in AF and VTE patients receiving anticoagulant therapy. This review summarises the literature involving original studies deriving bleeding risk scores and validation studies of these scores for stroke prevention in AF and treatment/prevention of VTE. To date, there are 10 bleeding risk scores, 6 for use in AF populations, three in VTE cohorts and 1 for mixed indications; they differ markedly in the number of, and risk factors for bleeding, and complexity. In conclusion, many clinical prediction tools to assess bleeding risk prior to starting OAC for either stroke prevention in AF or treatment of VTE are available and should be used in clinical practice to identify and manage modifiable risk factors.

Words: 174

Key words:

Bleeding risk score, atrial fibrillation, venous thromboembolism, anticoagulant therapy

Oral anticoagulants (OAC), including vitamin K antagonists (VKAs) and non-VKA oral anticoagulants (NOACs) are recommended to prevent thromboembolic events in patients with atrial fibrillation (AF) and for primary and secondary prevention of venous thromboembolism (VTE). However, OAC use is associated with an increased risk of major bleeding, the most serious of which is intracranial hemorrhage (ICH).¹ In clinical practice, estimating the bleeding risk of an individual patient is beneficial before starting an anticoagulant,² particularly to identify and manage modifiable bleeding risk factors. In addition bleeding risk assessment could help to identify those at higher bleeding risk for whom additional measures, such as more regular follow-up visits and good International Normalised Ratio (INR) management (for those on VKAs), providing information and/or practical measures to reduce the falls risk, and informing patients about high-risk activities¹ to reduce bleeding risk³ could be implemented. Bleeding risk scoring systems have been developed for use in AF and VTE patients to estimate bleeding risk and to help aid treatment decisions. These scores have been tested and validated worldwide in many cohorts of AF⁴⁻⁹ and VTE¹⁰⁻¹² to support physicians in assessing bleeding risks.¹³ More recently, the European Society of Cardiology (ESC) Guidelines on the management of AF summarised bleeding risks (into modifiable, potentially modifiable, non-modifiable and biomarker-based) and encouraged prompt attention to correct modifiable bleeding risks.¹⁴ This review summarises the original derivation and validation studies of these bleeding risk scores for stroke prevention in AF and treatment/prevention of VTE.

Methods

A literature search was performed in EMBASE (1974 to April 21 2016), EMBASE Classic (1947 to 1973), Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations and Ovid MEDLINE(R) (1946 to present). The search strategy included keywords and MeSH terms relating to bleeding risk score, major bleeding, bleeding, AF, VTE, warfarin, and oral anticoagulant individually and in combination. Primary published research articles on prospective or retrospective studies where bleeding risk scores were derived

and validated were included. Studies that did not provide comparative outcomes, information on follow up time, or were not published in the English language were excluded. A manual search of the reference lists was also performed for studies that were not retrieved by the database searches.

To date, 6 bleeding risk scores are available for use in AF populations,⁴⁻⁹ 3 in VTE populations¹⁰⁻¹² and 1 in a mixed indication population.¹⁵ Tables 1 to 4 summarises the main published prospective or retrospective studies where bleeding risk scores in patients with AF and VTE who are treated with warfarin were derived and/or validated.

Many different risk factors for bleeding are included in the various bleeding scores (**Table 1**), with the number of risk factors included ranging from three^{4,12} to 12.⁸ All bleeding scores^{4-12,15} included age, utilising different age ranges and cut-offs (ranging from > 50 years old to > 85 years old) to indicate greater risk of bleeding; four scores included age ≥ 75 years.^{5,6,8,11} After age, the most common bleeding risk factors included in the scores were as follows: (i) previous/remote bleeding (reported in 8 scores),^{4-9,11,15} (ii) renal disease (included in 7 scores),^{5-8,10,11,15} (iii) anemia (in 6 scores),^{5,6,8,9,11,15} malignancy (in 4 scores),^{8,10-12} liver disease,^{7,8,10} hypertension,⁶⁻⁸ stroke,^{7,8,15} combined antiplatelet therapy^{5,7,9} and alcohol excess,⁷⁻⁹ (all included in 3 scores), while two scores included diabetes mellitus,^{9,15} reduced platelet count^{8,10} and female sex.^{9,12}

Other risk factors included biomarkers (high-sensitivity troponin, cTnT-hs and growth differentiation factor-15, GDF-15), recent bleed, myocardial infarction, labile INR, excessive falls risk, genetic factors, active gastro-duodenal ulcer, Intensive Care Unit/Coronary Care Unit (ICU/CCU) admission, central venous catheter, rheumatic disease, male sex and clinically overt pulmonary embolism. Seven^{5,6,8-11,15} out of 10 studies reported patients with a history or diagnosis of cancer, as malignancy is associated with increased risk of bleeding,^{12,16} about 2.5%-30% had a history/diagnosis of cancer (**Table 2**). All 3¹⁰⁻¹² bleeding scores for VTE patients include history or current diagnosis of cancer in their bleeding score, as cancer is shown to increase the risk of thromboembolism.^{17,18} Interestingly the ORBIT⁵

score and the score developed by Ruiz-Gimenez et al¹¹ found almost 60% of their population had anemia, also a known risk factor for bleeding. In most bleeding scores^{4-7,9} for AF, hypertension was the most prevalent co-morbidity followed by congestive heart failure and diabetes.

Clinically overt PE was included as a risk factor for bleeding only in the VTE bleeding score by Ruiz-Gimenez et al.¹¹ The aim of the other 2 VTE scores^{10,12} was to develop a score with the ability to distinguish patients at low, mild or high risk of major bleeding during the first 90 days of OAC therapy. Kuijer et al¹² constructed the simplest bleeding risk prediction score, including only 3 clinical factors for patients at high risk of developing hemorrhagic complications at OAC inception, however this level of simplicity may miss other important features that could put patients at risk of bleeding.

Two bleeding risk scores, HEMORR₂HAGES⁸ and the ABC bleeding score,⁴ included factors that are not routinely available in daily clinical practice. HEMORR₂HAGES included genetic testing, although this was not available in their cohort, and the ABC score included 2 biomarkers, GDF-15 and cTnT-hs.

The mean/median age of study population in the derivation cohort ranged from 61¹⁵ to 80.2 years⁸ (**Table 2**). Almost half of the population in the derivation studies were female and only 3 studies^{5,6,15} reported ethnicity, which was predominantly White. Five out of 6^{4-7,9} studies from the AF cohorts reported hypertension as the most common co-morbid disease present in their population whereas one study from a VTE cohort¹⁰ and the mixed cohort¹⁵ reported kidney disease to be more prevalent in their patient population.

Table 3 presents the characteristics of derivation and validation studies of bleeding risk scores for AF and VTE. A prospective study design was used in 3^{4,5,7} out of 6 scores for AF populations and 2^{10,11} out of 3 studies in VTE populations. Two^{6,9} studies used a retrospective analysis and 1 study⁸ derived their score from the previous bleeding score available in AF, while a retrospective study design was used in 1¹² VTE study and 1 mixed population study.¹⁵

Most studies in AF patients had follow-up for at least 1 year except the first score developed by Shiremen et al⁹ which followed their patients for the first 90 days following hospital discharge after AF diagnosis. In contrast, bleeding scores derived from VTE cohorts usually followed up patients for 3 months, as the duration of treatment of DVT/PE with warfarin is typically between 3-12 months depending on the type of VTE.¹⁹ All studies derived their risk score using bleeding risk factors from large cohorts of patients ranging from 3456⁷ to almost 20,000⁹ patients, apart from 2 studies, by Kuijer and Landefeld et al. which only included 241 and 556 patients, respectively.^{12,15}

All bleeding risk scores stratified patients into 3 categories of bleeding risk (low, intermediate and high) except for the HAS-BLED score which initially categorised bleeding risk as high (score ≥ 3) and low-moderate risk (0-2)¹ and IMPROVE¹⁰ which categorized scores as ≥ 7 (increased risk of bleeding) and < 7 . These bleeding risk scores showed major bleeding rates ranging from 0.1%-3% in the low risk group and 4.9%-30% in the high risk group in the validation cohorts. **(Table 4)**

In terms of VTE prophylaxis, the most recent bleeding risk score developed to assess bleeding risks is the IMPROVE,¹⁰ and is perhaps the most comprehensive score by including more predictors of major bleeding (10 predictors), compared to the scores by Ruiz-Gimenez et al¹¹ (6 predictors) and Kuijer et al¹² (3 predictors).

The earliest bleeding score developed by Landefeld et al¹⁵ in 1989 derived 5 predictive factors of major bleeding in a mixed indication population. One of the original risk factors was AF but this was later removed when the score was validated, as its association with major bleeding was no longer significant in the validation cohort. Diabetes mellitus was substituted instead of AF as a new predictor of major bleeding.

The ability of the bleeding risk scores to predict bleeding risk has been validated both in cohorts to similar to ones where the score was derived (4 studies)^{6,7,9,11} and in independent validation cohorts (6 studies).^{4,5,8,10,12,20} In the validation and comparison study by Hijazi et al,⁴ the ABC score statistically outperformed the HAS-BLED and ORBIT scores in predicting major bleeding in both the derivation cohort

[0.68 (95% CI 0.66–0.70) vs. 0.61 (0.59–0.63) vs. 0.65 (0.62–0.67), respectively; ABC-bleeding vs. HAS-BLED $p<0.0001$ and ABC-bleeding vs. ORBIT $p=0.0008$] and the external validation cohort [0.71 (95% CI 0.68–0.73) vs. 0.62 (0.59–0.64) for HAS-BLED vs. 0.68 (0.65–0.70) for ORBIT; ABC-bleeding vs. HAS-BLED $p<0.0001$ and ABC-bleeding vs. ORBIT $p=0.0016$].⁴ Although the ABC score performed better than the HAS-BLED and ORBIT scores in this report, the complexity of the algorithm and inclusion of biomarkers which are not routinely performed in daily clinical practice, may make it difficult and more costly, for physicians to apply routinely.

One recent meta-analysis²¹ compared the diagnostic accuracy between HAS-BLED and HEMORR₂HAGES, ATRIA, CHADS₂ or CHA₂DS₂-VASc scores in anticoagulated patients with AF. The findings revealed that the HAS-BLED score performed better than the HEMORR₂HAGES and ATRIA bleeding scores, as well as being superior to CHADS₂ or CHA₂DS₂-VASc in predicting bleeding. Despite having better performance when compared to HEMORR₂HAGES, ATRIA and ORBIT, an additional advantage of the HAS-BLED score over the other five bleeding scores is the inclusion of quality of anticoagulation control (the 'L' acronym for labile INR or poor TTR<65%). Time in therapeutic (TTR) is a reflection of anticoagulation control in patients taking a VKA; a target TTR of $\geq 70\%$ is optimal for efficacy and safety.²²

In a post-hoc analysis evaluating the performance of HAS-BLED, ATRIA and ORBIT bleeding risk scores in the AMADEUS trial,²³ TTR was strongly correlated with clinically relevant bleeding events in patients using the ATRIA and ORBIT score, thus demonstrating that incorporating TTR in bleeding scores improves their ability to predict future bleeding events. Another comparison of four bleeding risk scores (HAS-BLED, ORBIT, ATRIA and HEMORR₂HAGES) in the SPORTIF cohort²⁴ also investigated whether the addition of 'labile INR' (TTR<65%) improved bleeding risk prediction (with the exception of the HAS-BLED score which already contains labile INR). Addition of 'labile INR' to the ORBIT, ATRIA and HEMORR₂HAGES bleeding risk scores, significantly improved the predictive performance of each

score for major bleeding [integrated discriminatory improvement (IDI) 0.0023, $p=0.0092$ vs. IDI 0.0020, $p=0.00014$ vs. IDI 0.0015, $p=0.0016$, respectively].²⁴

Apostolakis et al²⁵ compared the predictive performance of HAS-BLED with HEMORR₂HAGES and ATRIA in the AMADEUS trial and demonstrated that the HAS-BLED score performed better than HEMORR₂HAGES and ATRIA score in predicting any clinically relevant bleeding, with only the HAS-BLED score demonstrating significant improvement for intracranial hemorrhage.²⁵ In another ancillary analysis of the same trial,²⁶ the HAS-BLED score performed better than the ORBIT score in predicting any clinically relevant bleed in a non-oral anticoagulant (idraparinux).²⁶

More recently the predictive ability of the HAS-BLED score was also investigated in patients receiving NOAC therapy, with rivaroxaban, in a small retrospective case-control study;²⁷ the HAS-BLED score demonstrated modest diagnostic ability to predict major bleeding events although this was not statistically significant (c statistics=0.68; $p=0.07$).²⁷ Analyses have demonstrated that the HAS-BLED score not only performs well in predicting bleeding events in VKA treated patients with AF, it can also be used to predict bleeding events in non-VKA treated patients which is very useful as more AF patients are being treated with NOACs.²⁷

Whereas in the VTE cohorts, only the IMPROVE score has been shown to have good predictive ability for major [ROC 0.67 (95% CI, 0.57-0.77; $p=0.008$)] and clinically important bleeding [ROC 0.64 (95CI, 0.55-0.73; $p=0.0080$)] at 14 days when validated by Hostler et al in a cohort of 1688 hospitalised patients.²⁸

In conclusion, balancing individual risk of thromboembolic events and bleeding is complex but maximising the benefit of OAC while minimising bleeding risk, resulting in a net clinical benefit, should be undertaken in all patients receiving OAC. As reviewed here, there are many clinical prediction tools to assess bleeding risk prior to starting OAC for either stroke prevention in AF or treatment of VTE, which should be used in clinical practice to identify and manage modifiable risk factors.

Acknowledgments: None

Declaration of Interest

HZ: Received funding from Malaysian Ministry of Education and Universiti Teknologi MARA for her PhD study.

GYHL: Consultant for Bayer/Janssen, BMS/Pfizer, Biotronik, Medtronic, Boehringer Ingelheim, Microlife and Daiichi-Sankyo. Speaker for Bayer, BMS/Pfizer, Medtronic, Boehringer Ingelheim, Microlife, Roche and Daiichi-Sankyo

DAL: Investigator-initiated educational grants from Boehringer Ingelheim and Bristol Myers Squibb, speaker fees from Boehringer Ingelheim, Bayer, Bristol Myers Squibb, and Pfizer, consulting fees from Boehringer Ingelheim, Bayer, and Bristol Myers Squibb/Pfizer, and non-financial support from Boehringer Ingelheim and Bristol Myers Squibb, outside the submitted work.

Word count: 2013

1. Lip GH, Lane DA. Stroke prevention in atrial fibrillation: A systematic review. *JAMA* 2015;313:1950-1962.
2. Lip GY, Lane DA. Bleeding risk assessment in atrial fibrillation: observations on the use and misuse of bleeding risk scores. *J Thromb Haemost* 2016;14:1711-1714.
3. Lip GYH, Nieuwlaat R, Lane D, Pisters R, Crijns H. A simple, new risk factor based approach to refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation: The Euro Heart Survey on atrial fibrillation. *Chest* 2010;30:979-979.
4. Hijazi Z, Oldgren J, Lindbäck J, Alexander JH, Connolly SJ, Eikelboom JW, Ezekowitz MD, Held C, Hylek EM, Lopes RD, Siegbahn A, Yusuf S, Granger CB, Wallentin L. The novel biomarker-based ABC (age, biomarkers, clinical history)-bleeding risk score for patients with atrial fibrillation: a derivation and validation study. *Lancet* 2016;387:2302-2311.
5. O'Brien EC, Simon DN, Singer DE, Thomas LE, Hylek EM, Gersh B, Ansell JE, Kowey PR, Mahaffey KW, Chang P, Fonarow GC, Piccini JP, Peterson ED. The ORBIT bleeding score: a simple bedside score to assess bleeding risk in atrial fibrillation. *Eur Hear J* 2015;36:3258-3264.
6. Fang MC, Chang Y, Borowsky LH, Pomernacki NK, Udaltsova N, Singer DE. A new risk scheme to predict warfarin-associated hemorrhage. *J Am Coll Cardiol* 2011;58:395-401.
7. Pisters R, Lane DA, Nieuwlaat R, de Vos CB, Crijns HJGM, Lip GYH. A novel user-friendly score (HAS-BLED) to assess 1-year risk of major bleeding in patients with atrial fibrillation: the Euro Heart Survey. *Chest* 2010;138:1093-1100.
8. Gage BF, Yan Y, Milligan PE, Waterman AD, Culverhouse R, Rich MW, Radford MJ. Clinical classification schemes for predicting hemorrhage: results from the National Registry of Atrial Fibrillation (NRAF). *Am Heart J* 2006;151:713-719.
9. Shireman TI, Mahnken JD, Howard PA, Kresowik TF, Hou Q, Ellerbeck EF. Development of a contemporary bleeding risk model for elderly warfarin recipients. *Chest* 2006;130:1390-1396.
10. Decousus H, Tapson VF, Bergmann J-F, Chong BH, Froehlich JB, Kakkar AK, Merli GJ, Monreal M, Nakamura M, Pavanello R, Pini M, Piovella F, Spencer FA, Spyropoulos AC, Turpie AGG, Zotz RB,

FitzGerald G, Anderson FA. Factors at admission associated with bleeding risk in medical patients: Findings from the IMPROVE investigators. *Chest* 2011;139:69-79.

11. Ruiz-Gimenez N, Suarez C, Gonzalez R, Nieto JA, Todoli JA, Samperiz AL, Monreal M. Predictive variables for major bleeding events in patients presenting with documented acute venous thromboembolism. Findings from the RIETE Registry. *Thromb Haemost* 2008;100:26-31.

12. Kuijter PM, Hutten BA, Prins MH, Büller HR. Prediction of the risk of bleeding during anticoagulant treatment for venous thromboembolism. *Arch Intern Med* 1999;159:457-460.

13. Donzé J, Rodondi N, Waeber G, Monney P. Scores to predict major bleeding risk during oral anticoagulation therapy: a prospective validation study. *Am J Med* 2012;125:1095-1102.

14. Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B, Castella M, Diener H-C, Heidbuchel H, Hendriks J, Hindricks G, Manolis AS, Oldgren J, Popescu BA, Schotten U, Van Putte B, Vardas P, Agewall S, Camm J, Baron Esquivias G, Budts W, Caceris S, Casselman F, Coca A, De Caterina R, Deftereos S, Dobrev D, Ferro JM, Filippatos G, Fitzsimons D, Gorennek B, Guenoun M, Hohnloser SH, Kolh P, Lip GYH, Manolis A, McMurray J, Ponikowski P, Roseneck R, Ruschitzka F, Savelieva I, Sharma S, Suwalski P, Tamargo JL, Taylor CJ, Van Gelder IC, Voors AA, Windecker S, Zamorano JL, Zeppenfeld K. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J* 2016;37:2893-2962.

15. Landefeld CS, Goldman OL. Major bleeding in outpatients treated with warfarin: incidence and prediction by factors known at the start of outpatient therapy. *Am J Med* 1989;87:144-152.

16. Prandoni P, Lensing AWA, Piccioli A, Bernardi E, Simioni P, Girolami B, Marchiori A, Sabbion P, Prins MH, Noventa F, Girolami A. Recurrent venous thromboembolism and bleeding complications during anticoagulant treatment in patients with cancer and venous thrombosis. *Blood* 2002;100:3484-3488.

17. Lee AYY, Levine MN. Venous thromboembolism and cancer: risks and outcomes. *Circulation* 2003;107:I-17-I-21.

18. Ay C, Dunkler D, Marosi C, Chiriac A-L, Vormittag R, Simanek R, Quehenberger P, Zielinski C, Pabinger I. Prediction of venous thromboembolism in cancer patients. *Blood* 2010;116:5377-5382.

19. Guyatt GH, Akl EA, Crowther M, Gutterman DD, Schünemann HJ. Executive summary: Antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2012;141:7S-47S.
20. Beyth RJ, Quinn LM, Landefeld CS. Prospective evaluation of an index for predicting the risk of major bleeding in outpatients treated with warfarin. *Am J Med* 1998;105:91-99.
21. Zhu W, He W, Guo L, Wang X, Hong K. The HAS-BLED score for predicting major bleeding risk in anticoagulated patients with atrial fibrillation: a systematic review and meta-analysis. *Clin Cardiol* 2015;38:555-561.
22. Camm AJ, Lip GYH, De Caterina R, Savelieva I, Atar D, Hohnloser SH, Hindricks G, Kirchhof P, Bax JJ, Baumgartner H, Ceconi C, Dean V, Deaton C, Fagard R, Funck-Brentano C, Hasdai D, Hoes A, Knuuti J, Kolh P, McDonagh T, Moulin C, Popescu BA, Reiner Ž, Sechtem U, Sirnes PA, Tendera M, Torbicki A, Vahanian A, Windecker S, Vardas P, Al-Attar N, Alfieri O, Angelini A, Blömstrom-Lundqvist C, Colonna P, De Sutter J, Ernst S, Goette A, Gorenek B, Hatala R, Heidbüchel H, Heldal M, Kristensen SD, Le Heuzey J-Y, Mavrakakis H, Mont L, Filardi PP, Ponikowski P, Prendergast B, Rutten FH, Schotten U, Van Gelder IC, Verheugt FWA. 2012 focused update of the ESC Guidelines for the management of atrial fibrillation: an update of the 2010 ESC guideline for the management of atrial fibrillation. *Eur Heart J* 2012;33:2719-2747.
23. Senoo K, Proietti M, Lane DA, Lip GY. Evaluation of the HAS-BLED, ATRIA, and ORBIT bleeding risk scores in patients with atrial fibrillation taking warfarin. *Am J Med* 2016;129:600-607.
24. Proietti M, Senoo K, Lane DA, Lip GY. Major bleeding in patients with non-valvular atrial fibrillation: impact of time in therapeutic range on contemporary bleeding risk scores. *Sci Rep* 2016;6:24376.
25. Apostolakis S, Lane DA, Guo Y, Buller H, Lip GYH. Performance of the HEMORR₂HAGES, ATRIA, and HAS-BLED bleeding risk – Prediction scores in patients with atrial fibrillation undergoing anticoagulation. The AMADEUS (Evaluating the Use of SR34006 Compared to Warfarin or Acenocoumarol in Patients With Atrial Fibrillation) Study. *JACC* 2012;60:861-867.

26. Senoo K, Lip GY. Predictive abilities of the HAS-BLED and ORBIT bleeding risk scores in non-warfarin anticoagulated atrial fibrillation patients: an ancillary analysis from the AMADEUS trial. *Int J Cardiol* 2016;221:379-382.
27. Gorman EW, Perkel D, Dennis D, Yates J, Heidel RE, Wortham D. Validation of the HAS-BLED tool in atrial fibrillation patients receiving rivaroxaban. *J Atr Fibrillation* 2016;9:16-18.
28. Hostler DC, Marx ES, Moores LK, Petteys SK, Hostler JM, Mitchell JD, Holley PR, Collen JF, Foster BE, Holley AB. Validation of the international medical prevention registry on venous thromboembolism bleeding risk score. *Chest* 2016;149:372-379.

List of Tables

Table 1: Risk factors for bleeding included in each bleeding risk score

Table 2: Baseline patient characteristics of the derivation cohorts for each bleeding risk score

Table 3: Characteristics of the derivation and validation cohorts for each of the bleeding risk scores

Table 4: Risk factors, risk categories and bleeding events in the validation cohorts

Table 1: Risk factors for bleeding included in each bleeding risk score

Risk factor	ABC ⁴	ORBIT ⁵	ATRIA ⁶	HAS-BLED ⁷	HEMORR ₂ HA GES ⁸	Shireman ⁹	IMPROVE ¹⁰	Ruiz-Gimenez ¹¹	Kuijer ¹²	OBRI ¹⁵
Age ≥85							✓			
Age ≥75		✓	✓		✓			✓		
Age ≥70						✓				
Age ≥65				✓						✓
Age ≥60									✓	
Age ≥50	✓									
Biomarkers	✓									
Previous/remote bleed	✓	✓	✓	✓	✓	✓		✓		✓
Recent bleed						✓				
Anaemia		✓	✓		✓	✓		✓		✓
Renal disease		✓	✓	✓	✓		✓	✓		✓
Liver disease				✓	✓		✓			
Hypertension			✓	✓	✓					
Myocardial infarction										✓
Diabetes						✓				✓
Malignancy					✓		✓	✓	✓	
Stroke				✓	✓					✓
Combined antiplatelet therapy		✓		✓		✓				
Labile INR				✓						
Alcohol excess				✓	✓	✓				
Excessive fall risk					✓					
Genetic factors					✓					

Active gastro-duodenal ulcer							✓			
Bleeding 3 months before admission							✓			
Reduced platelet count					✓		✓			
ICU/CCU							✓			
Central venous catheter							✓			
Rheumatic disease							✓			
Male sex								✓		
Female sex						✓			✓	
Clinically overt PE								✓		
Total no. of risk factors/score	3	5	5	9	12	8	10	6	3	7

CCU = coronary care unit; ICU = intensive coronary care unit; INR = international normalised ratio; PE = pulmonary embolism

Table 2: Baseline patient characteristics of the derivation cohorts for each bleeding risk score

Patients, %	ABC⁴	ORBIT⁵†	ATRIA⁶†	HAS-BLED⁷‡	HEMORR₂ HAGES⁸‡	Shireman⁹	IMPROVE¹⁰	Ruiz-Gimenez¹¹	Kuijjer¹²	OBRI¹⁵
Number of patients	14,537	581	307	53	1604	19,875	10,866	314	241	556
Mean age (SD)/median (IQR)	70 (19-97)	78 (71–83)	-	73 (10)	80.2	88% ≥70 years	68.1 (51.8-78.9)	66 (17)	63 (17)	61 (14)
Sex (female)	36	46.1	37.4	53	57	52.5	50.6	45	46	54
Ethnicity	-	91.6	86.2	-	-	-	-	-	-	93
White	-	30.8	18.0	-	4.8	2.5	10.7	35	24	-
History/ diagnosis of cancer	-	57.5	18.8	-	8.5	7.5	-	56	-	-
Anaemia/abnormal Hb/Hct	-	89.3	64.7	74	0.4	72	38.7	-	-	-
Hypertension	87	33.7	22.1	23	-	29.6	-	-	-	8
Diabetes	25	44.9	-	45	-	59.8	10.8	7.6	-	-
CHF	31	20.5	0.5	29**	-	68.5**	13	-	-	4
MI	13	13.1	17.4	12	37.2	32.1	-	-	-	12
Prior stroke	19	15.5	12.1	17 [§]	-	11.9	-	5.4	-	10
GI bleed	-	48.4	5.9 [¶]	19	-	0.6 [#]	11.0 [¶]	-	-	18
eGFR 60ml/min	74.1 ml/min									
Antiplatelet	-	49.1	0.9	-	-	22.3	24.6	14	-	-
Warfarin	-	95	-	-	42.3	28.7	1.3*	-	-	-
NOACs	-	5.2	-	-	-	-	-	-	-	-

CAD = coronary artery disease; CHF = congestive heart failure; eGFR = estimated glomerular filtration rate; GI = gastrointestinal; Hb = hemoglobin; Hct = hematocrit; IQR = interquartile range; MI = myocardial infarction; NOACs = non-vitamin K antagonist oral anticoagulant; SD = standard deviation *anticoagulant discontinued; †population experiencing major bleed only; ‡ warfarin users only; § major bleed; ¶ renal failure; #hepatic/renal failure; ** CAD; - not reported

Table 3: Characteristics of the derivation and validation cohorts for each of the bleeding risk scores

Risk score	Year	Country	Derivation cohort				Validation cohort				Major bleed definition
			Indication for OAC	Study design	Subjects	Follow-up	Indication for OAC	Study design	Subjects	Follow-up	
ABC ⁴	2016	Sweden	AF	Prospective	14,537	1.7 years†	AF	Prospective	8468	1.9†	ISTH
ORBIT ⁵	2015	USA	AF	Prospective	7411	2 years	AF	Prospective	14,264	1.9†	ISTH
ATRIA ⁶	2011	USA	AF	Retrospective	6123	6 years	AF	Retrospective	3063	6 years	ISTH
HAS-BLED ⁷	2010	Europe	AF	Prospective	3456	1 year	AF	Prospective	3071	1 year	ISTH
HEMORR ₂ HAGES ⁸	2006	USA					AF	Retrospective	3791	3 years	Hospitalisation for bleeding
Shireman ⁹	2006	USA	AF	Retrospective	19,875	3 mth	AF	Retrospective	6470	3 mth	Hospitalisation for GI bleed or ICH
IMPROVE ^{10,28}	2011	USA	VTE	Retrospective or prospective	10,866	3 mth	VTE	Prospective	1668	18 mth	ISTH
Ruiz-Gimenez ¹¹	2008	Spain	VTE	Prospective	13,057	3 mth	VTE	Prospective	6572	3 mth	ISTH
Kuijjer ¹²	1999	The Netherlands	VTE	Retrospective	241	3 mth	VTE	Retrospective	780	3 mth	ISTH
OBRI ^{15,20}	1989	USA	Mixed‡	Retrospective	556	48 mth	Mixed‡	Prospective	264	48 mth	ISTH

AF = atrial fibrillation; DVT = deep vein thrombosis; GI = gastrointestinal; ICH = intracranial haemorrhage; ISTH = International Society of Thrombosis and Haemostasis; mth = months; PE = pulmonary embolism; TE = thromboembolism; TIA = transient ischaemic attack; VHD = valvular heart disease; VTE = venous thromboembolism, valvular heart surgery, mitral valve disease, AF, stroke, transient ischemic attack (TIA), pulmonary embolism (PE), deep vein thrombosis (DVT), and other thromboembolism

†median; ‡mixed indication included valvular heart disease, AF, stroke, transient ischemic attack, deep vein thrombosis, pulmonary embolism and other thromboembolism

Table 4: Risk factors, risk categories and bleeding events in the validation cohorts

Risk score	Risk factors (score for each factor)	Risk categories			Bleeding events in validation cohort (per 100 patient yrs)		
		Low	Intermediate	High	Low	Intermediate	High
ABC ⁴	Age(†); Biomarkers (†) (GDF-15 or cystatin C/CKD-EPI, cTnT-hs, & Hb); Previous bleed (†)	<1%	1-2%	>3%	0.62	1.67	4.87
ORBIT ⁵	Age ≥75 yrs (1); ↓Hb/Hct/anemia (2); Bleeding history (2); ↓ renal function (1); APT (1)	0-2	3	≥4	2.4*	4.7	8.1
ATRIA ⁶	Anemia (3); Severe renal disease (3); Age ≥75 yrs (2); Prior bleed (1); Hypertension (1)	0-3	4	5-10	0.83	2.41	5.32
HAS-BLED ⁷	↑SBP (1); Severe renal/hepatic disease (1 each); Stroke (1);Bleeding (1); Labile INR (1); Age >65 yrs (1); APT/NSAIDs (1); Alcohol excess (1)	0-1	2	≥3	1.02-1.13	1.88	≥3.74
HEMORR ₂ HAGES ⁸	Hepatic/renal disease (1); Ethanol abuse (1); Malignancy; Age >75 yrs (1); ↓Plt (1); Re-bleeding risk (2); ↑BP (1); Anemia (1); Genetic factors (1); ↑ falls risk (1); Stroke (1)	0-1	2-3	≥4	1.9-2.5	5.3-8.4	10.4-12.3
Shireman et al ⁹	Age ≥70 years (0.49); Female (0.31); Previous bleed (0.58); Recent bleed (0.62); Alcohol/drug abuse (0.71); DM (0.27); Anemia (0.86); APT (0.32)	≤1.07	>1.07/ <2.19	≥2.19	0.9% ^a	2.0% ^a	5.4% ^a
IMPROVE ^{10,28}	Active GI ulcer (4.5); Recent bleed (4); ↓Plt (4); Age ≥75 yrs (3.5); Hepatic/renal failure (2.5 each); ICU/CCU admission (2.5); CV catheter (2); Rheumatic disease (2); current cancer (2); Male (1)	<7	-	≥7	2.7%‡		6.5%‡
Ruiz-Gimenez et al ¹¹	Recent major bleed (2); ↑Creat (1.5); Anemia (1.5); Cancer (1); PE (1); Age >75 yrs (1)	0	1-4	>4	0.1% ^a	2.8% ^a	6.2% ^a
Kuijjer et al ¹²	Age≥60 yrs (1.6); Female (1.3); Malignancy (2.2)	0	1-3	>3	0.6% ^a	2% ^a	7% ^a
OBRI ^{15,20}	Age≥65 yrs (1); Previous stroke (1); Previous GI bleed (1); Recent MI/ anemia/DM/↑ creat (1)	0	1-2	3-4	3% ^b	8% ^b	30% ^b

APT = antiplatelet therapy; BP = blood pressure; CCU = coronary care unit; creat = creatinine; cTnT-hs = Troponin T; CV = central venous; DM = diabetes mellitus; GDF-15 = growth differentiation factor-15; GI = gastrointestinal; Hb = haemoglobin; Hct = hematocrit; ICU = intensive care unit; INR = international normalised ratio; MI = myocardial infarction; PE: pulmonary embolism; Plt = platelet count or function; SBP = systolic blood pressure; yrs = years

* bleeding event in original derivation cohort; ^a at 3 months; ^b at 12 months; ↓ reduced/decreased; ↑ elevated/increased; † score for each variable in ABC score is based on a nonogram (see reference⁴); ‡ clinically important bleeding: sum of major bleed and clinically relevant non-major